

- i. Active osteomyelitis of any bone or a substantiated history of osteomyelitis of any of the long bones of the extremities at any time.
- j. Filariasis, trypanosomiasis, amoebiasis or schistosomiasis.
- k. Hodgkin's disease.
- l. Uncinariasis, if more than mild.
- m. Malaria, chronic, if more than mild.
- n. Splenectomy for any cause, other than trauma or congenital hemolytic icterus.
- o. Leukemia.

## SECTION III

## HEIGHT, WEIGHT, AND CHEST MEASUREMENTS

	Paragraph
Table of standard and minimum acceptable measurements of height, weight, and circumference of chest.....	9
Directions for taking height and weight.....	10
General service.....	11
Limited service.....	12
Nonacceptable.....	13
General considerations.....	14

9. Table of standard and minimum acceptable measurements of height, weight, and circumference of chest.

Height (inches)	Standard		Minimum	
	Weight	Chest measurement at expiration	Weight	Chest measurement at expiration
	<i>Pounds</i>	<i>Inches</i>	<i>Pounds</i>	<i>Inches</i>
60.....	116	31¼	105	28¾
61.....	119	31½	107	29
62.....	122	31¾	109	29¼
63.....	125	32	111	29½
64.....	128	32¼	113	29¾
65.....	132	32½	115	30
66.....	136	32¾	117	30¼
67.....	140	33	121	30½
68.....	144	33¼	125	30¾
69.....	148	33½	129	31
70.....	152	33¾	133	31¼
71.....	156	34	137	31½
72.....	160	34¼	141	31¾
73.....	164	34½	145	32
74.....	168	34¾	149	32¼
75.....	172	35	153	32½
76.....	176	35¼	157	32¾
77.....	180	35½	161	33
78.....	184	35¾	165	33¼

10. Directions for taking height and weight.—*a.* The measuring rod should consist of a board at least 2 inches wide by 80 inches long, placed vertically, firmly fixed, with accurate graduations of ¼ inch between 58 inches and the top end. Obtain the height by placing